

References

- Bowden MG, Embry AE, Gregory CM. Physical therapy adjuvants to promote optimization of walking recovery after stroke. *Stroke Res Treat*. 2011; Epub 2011 Oct 12.
- English C, Hillier SL. Circuit class therapy for improving mobility after stroke. *Cochrane Database Syst Rev*. 2010; 7: CD007513.
- Ferrarin M, Rabuffetti M, Bacchini M, Casiraghi A, Castagna A, Pizzi A, Montesano A. Does gait analysis change clinical decision-making in poststroke patients? Results from a pragmatic prospective observational study. *Euro J Phys Rehabil Med*. 2015; 51: 171-84.
- Kwakkel G, Kollen B, Lindeman E. Understanding the pattern of functional recovery after stroke: Facts and theories. *Restor Neurol Neurosci*. 2004; 22: 281-99.
- Lang CE, Macdonald JR, Reisman DS, Boyd L, Jacobson Kimberley T, Schindler-Ivens SM, Hornby TG, Ross SA, Scheets PL. *Arch Phys Med Rehabil*. 2009; 90: 1692-8.
- Mayo NE, Wood-Dauphinee S, Ahmed S, et al. Disablement following stroke. *Disability & Rehabilitation*. 1999;21(5/6):258-268.
- Moore JL, Roth EJ, Killan C, Hornby TG. Locomotor training improves daily stepping activity and gait efficiency in individuals poststroke who have reached a “plateau” in recovery. *Stroke*. 2010; 41: 129-35.
- Mudge S, Barber A, Stott S. Circuit-based rehabilitation improves gait endurance but not usual walking activity in chronic stroke: A randomized controlled trial. *Arch Phys Med Rehabil*. 2009; 90: 1989-96.
- Observational Gait Analysis Handbook. The Pathokinsiology Service and the Physical Therapy Department, Rancho Los Amigos National Rehabilitation Center, Downey, CA. 2001.
- Page SJ, Gater DR, Bach-y-Rita P. Reconsidering the motor recovery plateau in stroke rehabilitation. *Arch Phys Med Rehabil*. 2004; 85: 1377-81.
- Pak S, Patten C. Strengthening to promote functional recovery poststroke: An evidence-based review. *Top Stroke Rehabil*. 2008; 15: 177-99.
- Perry J, Burnfield J. *Gait Analysis. Normal and Pathological Function*. Slack Inc, Thorofare, NJ. 2nd Ed. 2010

Seale J, Utsey C. Physical Therapists' Clinical Decision Making in Patients with Gait Dysfunction from Hemiplegia, 2012; from unpublished data.

Teasell RW, Bhogal SK, Foley NC, Speechley MR. Gait retraining post stroke. *Top Stroke Rehabil.* 2003; 10: 34-65.

Teasell RW, Murie Fernandez M, McIntyre A, Mehta S. Rethinking the continuum of stroke rehabilitation. *Top Stroke Rehabil.* 2014; 95: 595-6

Weiss A, Suzuki T, Bean J, Fielding RA. High intensity strength training improves strength and functional performance after stroke. *Am J Phys Med Rehabil.* 2000; 79: 369-76.

Wevers L, van de Port I, Vermue M, Mead G, Kwakkel G. Effects of task-oriented circuit class training on walking competency after stroke. *Stroke.* 2009; 40: 2450-59.