

References: Elevating the Intensity Stroke Rehabilitation: Locomotor, Functional Mobility and Exercise Applications

- 1) Adherence to accelerometry measurement of community ambulation poststroke. *Phys Ther.* 2014 Jan;94(1):101-10. Barak S1, Wu SS,
- 2) Rose DK1, Behrman AL, Nadeau SE, Martin AD, Schofield RS, Tilson JK, Cen SY, Lu X, Wu SS, Azen, S, Duncan PW; Does exercise tolerance testing at 60 days poststroke predict rehabilitation performance? *Arch Phys Med Rehabil.* 2013 Jul;94(7):1223-9. doi: 10.1016/j.apmr.2013.01.031.
- 3) Nadeau SE1, Wu SS, Dobkin BH, Azen SP, Rose DK, Tilson JK, Cen SY, Duncan PW; Effects of task-specific and impairment-based training compared with usual care on functional walking ability after inpatient stroke rehabilitation: LEAPS Trial. *Neurorehabil Neural Repair.* 2013 May;27(4):370-80.
- 4) Wulf G, Chiviacowski S, Lewthwaite R. Altering mindset can enhance motor learning in older adults. *Psychology and Aging.* 2012;27(1):14-21.
- 5) Boonsinsukh R, Panichareon L, Phansuwan-Pujito P. Light touch cue through a cane improves pelvic stability during walking in stroke. *Arch Phys Med Rehabil.* 2009;90:919-926.
- 6) Yen CL, Wang RY, Liao KK, Huang CC, Yang YR. Gait training induced change in corticomotor excitability in patients with chronic stroke. *Neurorehabilitation and Neural Repair.* 2008;22(1):22-30.
- 7) Bang DH, Shin WS, Kim SY, Choi JD. The effects of action observational training on walking ability in chronic stroke patients: a double-blind randomized controlled trial. *Clin Rehabil.* 2013 Oct 2.
- 8) Pons JL, Moreno JC, Torricelli D, Taylor JS. Principles of human locomotion: A review. *Conf Proc IEEE Eng Med Biol Soc.* 2013 Jul;2013:6941-4.
- 9) Bujanda ED, Nadeau S, Bourbonnais D, Dickstein R. Associations between lower limb impairments, locomotor capacities and kinematic variables in the frontal plane during walking in adults with chronic stroke. *J Rehabil Me.* 2003;35:259-264.