

Evidence-Based Interventions for Dementia Behaviors

- Achterberg, W. P., Pieper, M. J. C., van Dalen-Kok, A. H., de Waal, B. S., Lautenbacher, S., . . . Corbett, A. (2013). Pain management in patients with dementia. *Clinical Interventions in Aging, 8*, 1471-1482.
- Ahn, H. & Horgas, A. (2013). The relationship between pain and disruptive behaviors in nursing home residents with dementia. *BMC Geriatrics, 13*:14.
- Alessi, C., Yoon, E., Schnelle, F., Al-Samarrai, N., & Cruise, P. (1999). A randomized trial of a combined physical activity and environmental intervention in nursing home residents: do sleep and agitation improve? *Journal of the American Geriatric Society, 47*(7), 784-791.
- Alessi, C., Martin, J., Webber, A., et al. (2005). Randomized, controlled trial of a nonpharmacological intervention to improve abnormal sleep/wake patterns in nursing home residents. *Journal of the American Geriatric Society, 53*(5), 803-810.
- Algate, D., Beck, C., Kolanowski, A., Whall, A., Berent, S., Rickards, K., & Beattie, E. (1996). Need-driven dementia-compromised behavior: An alternative view of disruptive behavior. *American Journal of Alzheimer's Disease, 11*(6), 10-19.
- Allen, C. K., & Blue, T. (1998). Cognitive disabilities model: How to make clinical judgments. In N. Katz (Ed.), *Cognition and occupation in rehabilitation: Cognitive models for intervention in occupational therapy* (pp. 225-280). Bethesda, MD: American Occupational Therapy Association.
- Allen-Burge, R., Stevens, A., & Burgio, L. (1999). Effective behavioral interventions for decreasing dementia-related challenging behavior in nursing homes. *International Journal of Geriatric Psychiatry, 14*(3), 212-28.
- Amella, E. (1999). Factors influencing the proportion of food consumed by nursing home residents with dementia. *Journal of the American Geriatrics Society, 47*(7), 879-85.
- Beck, C. (1997). Improving dressing behavior in cognitively impaired nursing home residents. *Nursing Research, 46*(3), 126-132.
- Beck, C., et al. (2002). Effects of behavioral interventions on disruptive behavior and affect in demented nursing home residents. *Nursing Research, 51*(4), 219-28.
- Boustani, M., Callahan, C. M., Unverzagt, F. W., Austrom, M. G., Perkins, A. J., . . . Hendrie, H. C. (2005). Implementing a screening and diagnosis program for dementia in primary care. *Journal of General Internal Medicine, 20*, 572-577.
- Boustani, M., Zimmerman, S., Williams, C. S., Gruber-Baldini, A. L., Watson, L., . . . Sloane, P. D. (2005). Characteristics associated with behavioral symptoms related to dementia in long-term care residents. *The Gerontologist, 45*(special issue 1), 56-61.
- Braben, I. (1992). A Song for Mrs. Smith. *Nursing Times, 88*(41), 54.

- Buffington, A. L. H., Lipski, D. M., & Westfall, E. (2013). Dementia: An evidence-based review of common presentations and family-based interventions. *The Journal of the American Osteopathic Association*, 113(10), 768-775.
- Buhr, G. & White, H. (2007). Difficult behaviors in long-term care patients with dementia. *Journal of the American Medical Directors Association*, 8(3 suppl 2):e101e113.
- Burgener, S. & Twigg, P. (2002). Interventions for Persons with Irreversible Dementia. *Annual Review of Nursing Research*, 20, 89–124.
- Burgio, L., Stevens, A., Burgio, K., Roth, D., Paul, P., & Gerstle, J. (2002). Teaching and maintaining behavior management skills in the nursing home. *Gerontologist*, 42(4), 487-496.
- Cameron, M., Lonergan, E., & Lee, H. (2009). Transcutaneous Electrical Nerve Stimulation (TENS) for dementia. *Cochrane Database of Systematic Reviews*, (3).
- Casby, J. & Holm, M. (1994). The effect of music on repetitive disruptive vocalizations of persons with dementia. *American Journal of Occupational Therapy*, 48(10), 883-889.
- Cera, M. L., Ortiz, K. Z., Bertolucci, P. H. F., & Minett, T. S. C. (2013). Speech and orofacial apraxias in Alzheimer's disease. *International Psychogeriatrics*, 25(10), 1679-1685.
- Chung, J. & Lai, C. (2009). Snoezelen for dementia. *Cochrane Database of Systematic Reviews*, (3).
- Clark, M., Lipe, A., & Bilbrey, M. (1998). Use of Music to Decrease Aggressive Behaviors in People with Dementia. *Journal of Gerontological Nursing*, 24(7), 10-7.
- Cohen-Mansfield, J., & Werner, P. (1998). The effects of an enhanced environment on nursing home residents who pace. *Gerontologist*, 38, 199–208.
- Cohen-Mansfield, J. (2013). Nonpharmacologic treatment of behavioral disorders in dementia. *Current Treatment Options in Neurology*, 15(6), 765-85.
- Corcoran, M. A. (2001). Dementia. In B. R. Bonder & M. B. Wagner (Eds.), *Functional performance in older adults* (2nd ed., pp. 287–304). Philadelphia, F.A. Davis.
- Cowan, N. (2008). What are the differences between long-term, short-term, and working memory? *Progress in Brain Research*, 169, 323-338.
- Coyne, M. L., & Hoskins, L. (1997). Improving eating behaviors in dementia using behavioral strategies. *Clinical Nursing Research*, 6, 275–290.
- Crispi, E. L., & Heitner, G. (2002). An activity-based intervention for caregivers and residents with dementia in nursing homes. *Activities, Adaptation & Aging*, 26, 61–72.

- Cuddy, L., & Duffin, J. (2005). Music, memory, and Alzheimer's disease: is music recognition spared in dementia, and how can it be assessed? *Medical Hypotheses*, 64(2), 229-235.
- Day, K., Carreon, D., & Stump, C. (2000). The Therapeutic Design of Environments for People with Dementia: A Review of the Empirical Research. *The Gerontologist*, 40(4), 397-416.
- Dello Russo, C., DiGiulio, P., Brunelli, C., Dimonte, V., Villani, D., . . . Toscani, F. (2008). Validation of the Italian version of the Discomfort Scale – Dementia of Alzheimer Type. *Journal of Advanced Nursing*, 64(3), 298-304.
- Detweiler, M. B., Kim, K. Y., & Taylor, B. Y. (2005). Focused supervision of high-risk fall dementia patients: A simple method to reduce fall incidence and severity. *American Journal of Alzheimer's Disease and Other Dementias*, 20, 97–104. doi: 10.1177/153331750502000205
- de Werd, M. M.E., Boelen, D., Rikkert, M. G. M. O., & Kessels, R. P. C. (2013). Errorless learning of everyday tasks in people with dementia. *Clinical Interventions in Aging*, 8, 1177-1190.
- Dickinson, J. I., McLain-Kark, J., & Marshall-Baker, A. (1995). The effects of visual barriers on exiting behavior in a dementia care unit. *Gerontologist*, 35, 127–130.
- Dobbs, D., Munn, J., Zimmerman, S., Boustani, M., Williams, C. S., . . . Reed, P. S. (2005). Characteristics associated with lower activity involvement in long-term care residents with dementia. *The Gerontologist*, 45(special issue 1), 81-86.
- Donovan, C., & Dupuis, M. (2000). Specialized care unit: Family and staff perceptions of significant elements. *Geriatric Nursing*, 21, 30–33.
- Doody, R., et al. (2001). Practice Parameter (an evidence-based review): Report of the Quality Standards Subcommittee to the American Academy of Neurology. *Neurology*, 56(9), 1154-1166.
- Etgen, T., Sander, D., Bickel, H., & Förstl, H. (2011). Mild cognitive impairment and dementia: The importance of modifiable risk factors. *Deutsches Ärzteblatt International*, 108(44), 743-750.
- Farless, L. B., & Ritchie, C. S. (2012). Challenges of pain management in long-term care. *Annals of Long-Term Care*, 20(5). Retrieved from <http://www.annalsoflongtermcare.com/article/challenges-pain-management-long-term-care>
- Forbes, D., Morgan, D., Bangma, J., Peacock, S., & Adamson, J. (2009). Light Therapy for Managing Sleep, Behaviour, and Mood Disturbances in Dementia. *Cochrane Database of Systematic Reviews*, (3).
- Galimberti, S., & Scarpini, E. (2013). Dementia: Not only Alzheimer's disease, an eye on frontotemporal dementia. *Acta Médica Portuguesa*, 26(4), 299-300.

- Gallini, A., Gardette, V., Vellas, B., Lapeyre-Mestre, M., Andrieu, S., & Brefel-Courbon, C. (2013). Persistent use of analgesic medications in mild-to-moderate Alzheimer's disease. *Drugs & Aging*, 30(6), 439-445.
- Gasecki, D., Kwarciany, M., Nyka, W., & Narkiewicz, K. (2013). Hypertension, brain damage and cognitive decline. *Current Hypertension Reports*, 15(6), 547-558.
- Gitlin, L. N., Winter, L., Burke, J., Chernett, N., Dennis, M. P., & Hauck, W. W. (2008). Tailored activities to manage neuropsychiatric behaviors in persons with dementia and reduce caregiver burden: A randomized pilot study. *American Journal of Geriatric Psychiatry*, 16, 229–239.
- Gleeson, M., & Timmins, F. (2004). The use of touch to enhance nursing care of older person in long-term mental health care facilities. *Journal of Psychiatric & Mental Health Nursing*, 11(5), 541-545.
- Goldfeld, K. S., Grabowski, D. C., Caudry, D. J., & Mitchell, S. L. (2013). Health insurance status and the care of nursing home residents with advanced dementia. *JAMA Internal Medicine*, 173(22).
- Gotell, E., Brown, S., & Ekman, S. (2003). Influence of caregiver singing and background music on posture, movement, and sensory awareness in dementia care. *International Psychogeriatrics*, 15, 411–430.
- Greer, K., Pustay, K., Zaun, T., & Coppens, P. (2001). A comparison of the effects of toys versus live animals on the communication of patients with dementia of the Alzheimer's type. *Clinical Gerontologist: The Journal of Aging and Mental Health*, 24(3-4), 157-182.
- Gregorio, G., Ramirez, D., & Ribera, C. (2003). Dementia and Nutrition. Intervention Study in Institutionalized Patients with Alzheimer Disease. *Journal of Nutrition, Health & Aging*, 7(5), 304-308.
- Hall, G., & Buckwalter, K. (1987). Progressively lowered stress threshold: A conceptual model of care of adults with Alzheimer's disease. *Archives of Psychiatric Nursing*, 1, 399–406.
- Hansen, N., Jorgensen, T., & Ortenblad, L. (2009). Massage and touch for dementia. *Cochrane Database of Systematic Reviews*, (3).
- Harada, C. N., Natelson Love, M. C., & Triebel, K. (2013). Normal cognitive aging. *Clinical Geriatric Medicine*, 29(4), 737-752.
- Hauer, K., Becker, C., Lindemann, U., & Beyer, N. (2006). Effectiveness of physical training on motor performance and fall prevention in cognitively impaired older persons: A systematic review. *American Journal of Physical Medicine and Rehabilitation*, 85, 847–857. doi: 10.1097/01.phm.0000228539.99682.32

- Heyn, P. (2003). The effect of a multisensory exercise program on engagement, behavior, and selected physiological indexes in persons with dementia. *American Journal of Alzheimer's Disease & Other Dementias*, 18, 247–251.
- Hoffman, F., van den Bussche, H., Wiese, B., Glaeske, G., & Kaduszkiewicz, H. (2014). Diagnoses indicating pain and analgesic drug prescription in patients with dementia: A comparison to age- and sex-matched controls. *BMC Geriatrics*, 14:20.
- Holmes, C., Hopkins, V., Hensford, C., MacLaughlin, V., Wilkinson, D., & Rosenvinge, H. (2002). Lavender oil as a treatment for agitated behaviour in severe dementia: A placebo controlled study. *International Journal of Geriatric Psychiatry*, 17, 305–308.
- Holt, F., Birks, T., Thorgrimsen, L., Spector, A., Wiles, A., & Orrell, M. (2009). Aroma therapy for dementia. *Cochrane Database of Systematic Reviews*, (3).
- Husebo, B. S., Ballard, C., Sandvik, R., Nilsen, O. B., & Aarsland, D. (2011). Efficacy of treating pain to reduce behavioral disturbances in residents of nursing homes with dementia: Cluster randomized clinical trial. *BMJ*, 343:d4065.
- International Psychogeriatric Association. Behavior and psychological symptoms of dementia (BPSD) educational pack. www.ipaonline.net/pdfs/1BPSDfinal.pdf [2]. Published 2002. Accessed November 10, 2015.
- Johnson, M. (2008). *Electrotherapy: Evidence-Based Practice*: Ed. Watson. T. Elsevier.
- Kapusta, P., Regier, L., Bareham, J., & Jensen, B. (2011). Behaviour management in dementia. *Canadian Family Physician*, 57, 1420-1422.
- Kincaid, C., & Peacock, J. R. (2003). Effect of a wall mural on decreasing four types of door-testing behaviors. *Journal of Applied Gerontology*, 22, 76–88.
- Kirk-Sanchez, N. J., & McGough, E. L. (2014). Physical exercise and cognitive performance in the elderly: Current perspectives. *Clinical Interventions in Aging*, 9, 51-62.
- Kitwood, T. & Bredin, K. (1992). Towards a theory of dementia care: Personhood and well-being. *Ageing and Society*, 12(3), 269-287.
- Kolanowski, A. (2001). Capturing interests: Therapeutic recreation activities for persons with dementia. *Therapeutic Recreation Journal*, 35, 220–235.
- Koss, E., & Gilmore, G. C. (1998). Environmental interventions and functional ability of AD patients. In B. Vellas, J. Fitten, & G. Frisoni (Eds.), *Research and practice in Alzheimer's disease* (pp. 185–193). New York: Springer.

- Kovach, C. R., Noonan, P. E., Schlidt, A. M., Reynolds, S., & Wells, T. (2006). The Serial Trial Intervention: an innovative approach to meeting needs of people with dementia. *Journal of Gerontological Nursing*, 32(4), 18-36.
- Kovach, C. R., Simpson, M. R., Joosse, L., Noonan, P. E., Reynolds, S. A., . . . Raff, H. (2012). Comparison of the effectiveness of two protocols for treating nursing home residents with advanced dementia. *Research in Gerontological Nursing*, 5(4), 251-263.
- Larson, E. B. (2010). Prospects for delaying the rising tide of worldwide, late-life dementias. *International Psychogeriatrics*, 22(8), 1196-1202.
- Leong, I. Y., Chong, M. S., & Gibson, S. J. (2006). The use of a self-reported pain measure, a nurse-reported pain measure and the PAINAD in nursing home residents with moderate and severe dementia: A validation study. *Age and Ageing*, 35, 252-256.
- Lewis, S. C. (2003). Elder care in occupational therapy. Thorofare, NJ: Slack.
- Livingston, G., Johnston, K., Katona, C., Paton, J., & Lyketsos, C. (2005). Old Age Task Force of the World Federation of Biological Psychiatry: Systematic review of psychological approaches to the management of neuropsychiatric symptoms of dementia. *American Journal of Psychiatry*, 162(11), 996-2021.
- Logsdon, R., McCurry, S., & Teri, L. Evidence-based psychological treatments for disruptive behaviors in individuals with dementia. *Psychology of Aging*, 22, 28-36.
- Martindale, B. (2008). Effect of animal-assisted therapy on engagement of rural nursing home residents. *American Journal of Recreation Therapy*, 7(4), 45-53.
- McCabe, B., Baun, M., Speich, D., & Agrawal, S. (2002). Resident dog in the Alzheimer's special care unit. *Western Journal of Nursing Residency*, 24(6), 684-696.
- McCurry, S., Gibbons, L., Logsdon, R., et al. (2005). Nighttime insomnia treatment and education for Alzheimer's disease: a randomized, controlled trial. *Journal of the American Geriatric Society*, 53(5):793-802.
- Mitchell, S. L., Teno, J. M., Kiely, D. K., Shaffer, M. L., Jones, R. N., . . . Hamel, M. B. (2009). The clinical course of advanced dementia. *New England Journal of Medicine*, 361(16), 1529-1538.
- Namazi, K. H., Rosner, T. T., & Calkins, M. P. (1989). Visual barriers to prevent ambulatory Alzheimer's patients from exiting through an emergency door. *Gerontologist*, 29, 699-702.
- Namazi, K. H., & Johnson, B. D. (1992). The effects of environmental barriers on the attention span of Alzheimer's disease patients. *American Journal of Alzheimer's Care and Related Disorders & Research*, 7, 9-15.

- Nygård, L., & Öhman, A. (2002). Managing changes in everyday occupations: The experience of persons with Alzheimer's disease. *OTJR: Occupation, Participation, and Health*, 22, 70–81.
- Neal, M., & Barton Wright, P. (2009). Validation therapy for dementia. *Cochrane Database of Systematic Reviews*, (3).
- O'Connor, D., Ames, D., Gardner, B., & King, M. (2009). Psychosocial treatments of behavior symptoms in dementia: a systematic review of reports meeting quality standards. *International Psychogeriatrician*, 21(2), 225-240.
- Oliver, D., Connelly, J. B., Victor, C. R., Shaw, F. E., Whitehead, A., Genc, Y., et al. (2007). Strategies to prevent falls and fractures in hospitals and effect of cognitive impairment: Systematic review and meta-analyses. *British Medical Journal*, 334, 82. doi: 10.1136/bmj.39049.706493.55
- Orsulic-Jeras, S., Judge, K., & Camp, C. (2000). Montessori-based activities for long-term care residents with advanced dementia: effects on engagement and affect. *The Gerontologist*, 40, 107–111.
- Padilla, R. (2011). Effectiveness of interventions designed to modify the activity demands of the occupations of self-care and leisure for people With Alzheimer's Disease and related dementias. *American Journal of Occupational Therapy*, 65(5), 523-531.
- Patel, K. V., Guralnik, J. M., Dansie, E. J., & Turk, D. C. (2013). Prevalence and impact of pain among older adults in the United States: Findings from the 2011 National Health and Aging Trends study. *Pain*, 154(12).
- Pomeroy, V. M. (1993). The effect of physiotherapy input on mobility skills of elderly people with severe dementing illness. *Clinical Rehabilitation*, 7, 163–170.
- Pool, J. (2001). Making contact: An activity-based model of care. *Journal of Dementia Care*, 9, 24–26.
- Ragneskog, H., Brane, G., Karlsson, I., & Kihlgren, M. (1996). Influence of Dinner Music on Food Intake and Symptoms Common in Dementia. *Scandinavian Journal of Caring Sciences*, 10, 11-17.
- Rogers, J. C., Holm, M. B., Burgio, L. D., Granieri, E., Hsu, C., & Hardin, J. M. (1999). Improving morning care routines of nursing home residents with dementia. *Journal of the American Geriatrics Society*, 47, 1049–1057.
- Satoh, M., Ogawa, J., Tokita, T., Nakaguchi, N., Nakao, K., . . . Tomimoto, H. (2014). The effects of physical exercise with music on cognitive function of elderly people: Mihama-Kiho project. *PLOS One*, 9(4), e95230.
- Swartz, K., Hantz, E., Crummer, G. Walton, J., & Frisina, R. (1989). Does the melody linger on? Music cognition in Alzheimer's disease. *Seminar on Neurology*, 9(2), 152-158.

- Shaw, F., Bond, J., Richardson, D., Dawson, P., Steen I., McKeith, I., et al. (2003). Multifactorial intervention after a fall in older people with cognitive impairment and dementia presenting to the accident and emergency department: Randomized controlled trial. *British Medical Journal*, 326, 73–90.
- Sherratt, K., Thornton, A., & Hatton, C. (2004). Music interventions for people with dementia: A review of the literature. *Aging & Mental Health*, 8(1), 3–12.
- Skovdahl, K., Kihlgren, A. L., & Kihlgren, M. (2003). Dementia and aggressiveness: Video recorded morning care from different care units. *Journal of Clinical Nursing*, 12, 888–898.
- Skjerve, A., Bjorvatn B., & Holsten, F. (2004). Light therapy for behavioural and psychological symptoms of dementia. *International Journal of Geriatric Psychiatry*, 19, 516–522.
- Stegemöller, E. L., Wilson, J. P. Hazamy, A., Shelley, M. C., Okun, M. S., . . . Hass, C. J. (2014). Associations between cognitive and gait performance during single- and dual-task walking in people with Parkinson disease. *Physical Therapy*, 94(6), 757-766.
- Takai, Y., Yamamoto-Mitani, N., Okamoto, Y., Koyama, K., & Honda, A. (2010). Literature review of pain prevalence among older residents of nursing homes. *Pain Management Nursing*, 11(4), 209-223.
- Telenius, E. W., Engedal, K., & Bergland, A. (2013). Physical performance and quality of life of nursing home residents with mild and moderate dementia. *International Journal of Environmental Research and Public Health*, 10, 6672-6686.
- Teno, J. M., Dosa, D., Rochon, T., Casey, V., & Mor, V. (2008). Development of a brief survey to measure nursing home residents' perceptions of pain management. *Journal of Pain and Symptom Management*, 36(6), 572-583.
- Thorgrimsen, L., Spector, A., Wiles, A., & Orrell, M. (2003). Aromatherapy for dementia. Cochrane Database of Systematic Reviews, Issue 3. Art. No.: CD003150. DOI: 10.1002/14651858.CD003150.
- Volicer, L., & van der Steen, J. T., (2014). Outcome measures for dementia in the advanced stage and at the end of life. *Advances in Geriatrics*, 2014, Article ID 346485.
- Ward, J. D. (2003). Psychiatric diagnoses and related intervention issues. In E. B. Crepeau, E. S. Cohn, & B. A. B. Schell (Eds.), *Willard & Spackman's occupational therapy* (10th ed., pp. 835–859). Philadelphia: Lippincott Williams & Wilkins.
- Warren, J. D., Rohrer, J. D., & Rossor, M. N. (2013). Frontotemporal dementia. *BMJ*, 347, f4827.
- Watson, R., & Green, S. (2006). Feeding and dementia: a systematic literature review. *Journal of Advanced Nursing*, 54, 86-93.

- Weina, P., Zhao, H., Zhishun, L. & Shi, W. (2009). Acupuncture for vascular dementia. *Cochrane Database of Systematic Reviews*, (2).
- Whall, A. L, Black, M. E., Groh, C. J., Yankou, D. J., Kupferschmid, B. J., & Foster, M. L. (1997). The effect of natural environments upon agitation and aggression in late stage dementia patients. *American Journal of Alzheimer's Diseases*, 12, 216–220.
- Woods, B., Spector, A., Jones, C., Orrell, M., & Davies, S. (2009). Reminiscence therapy for dementia. *Cochrane Database of Systematic Reviews*, (3).
- Yang, J., Mann, W. C., Nochajski, S., & Tomita, M. R. (1997). Use of assistive devices among elders with cognitive impairment: A follow-up study. *Topics in Geriatric Rehabilitation*, 13, 13–31.
- Zetteler, J. (2008). Effectiveness of simulated presence therapy for individuals with dementia: a systematic review and meta-analysis. *Aging and Mental Health*, 12(6), 779-785.