

References

- Almurdhi M, Reeves N, Bowling F et al. Reduced lower limb muscle strength and volume in patients with type 2 diabetes in relation to neuropathy, intramuscular fat and vitamin D levels. *Diabetes Care*. 2016; 39: 441-447.
- Centers for Disease Control and Prevention. *National Diabetes Statistics Report, 2017*. Atlanta, GA: Centers for Disease Control and Prevention, US Department of Health and Human Services.
- American Diabetes Association. Summary of Revisions of Standards of Medical Care in Diabetes-2019. Abridged for Primary Care Providers. *Clinical Diabetes*. 2019: 37:11-34.
- National Institute of Diabetes and Digestive and Kidney Disease. Eating Right for Kidney Health. 2016. <https://www.niddk.nih.gov/health-information/kidney-disease/chronic-kidney-disease-ckd/eating-nutrition>
- Snowling N, Hopkin W. Effects of different modes of exercise training on glucose control and risk for complications in type 2 diabetic patients: A meta-analysis. *Diabetes Care*. 2006: 29; 2518-2527.
- Coldberg S, Sigal R, Yardley J et al. Physical Activity/exercise and diabetes: A position statement of the American Diabetes Association. *Diabetes Care*. 2016: 39; 2067-2079.
- Holman R, Paul S, Bethel A et al. 10-year follow-up of intensive glucose control in type 2 diabetes. *N Engl J Med*. 2008: 359; 1577-1589.
- Power S, Howley E. Exercise Physiology: Theory and Application to Fitness and Performance. 10th ed. McGraw Hill Education, 2015.
- American College of Sport Medicine. ACSM's Guidelines for Exercise Testing and Prescription. 10th ed. Wolter Kluwer Health, 2016.
- Duviver B, Schaper N, Hesselink M et al. Breaking sitting with light activity vs structure exercise: a randomized crossover study demonstrating benefits of glycaemic control and insulin sensitivity in type 2 diabetes. *Diabetologia*. 2017: 60; 490-498.
- Biswas A, Oh P, Faulkner G. Sedentary time and its association with risk for disease, mortality and hospitalization in adults: a systematic literature review and meta-analysis. *Ann Intern Med*. 2015: 162; 123-132.
- Cassidy S, Thoma C, Houghton D. High-intensity interval training: a review of its impact on glucose control and cardiometabolic health. *Diabetologia*. 2017: 60; 7-23.
- Batacan R, Duncan M, Dalbo V et al. Effects of high-intensity interval training on cardiometabolic health: a systematic review of and meta-analysis of intervention studies. *Br J Sports Med*. 2016: 16; 1-12.
- American Heart Association. Hypertension Guidelines Resources. 2018
<https://www.heart.org/en/health-topics/high-blood-pressure/high-blood-pressure-toolkit-resources>

Kluding P, Pasnoom M, Singh R et al. The effects of exercise on neuropathic symptoms, nerve fiber function, and cutaneous innervation in people with diabetic peripheral neuropathy. *J Diabetes Complications*. 2012; 5: 424-429.

Umpierre D, Riberio P, Kramer C et al. Physical activity advice or structured exercise training and association with HbA1c levels in type 2 diabetes: a systematic review and meta-analysis. *JAMA*. 17: 1790-1799.

Phas K, Jia F, Kamper S. Effects of regular exercise training in adults with chronic kidney disease (PEDro synthesis update). *Br J Sports Med*. 2016. 50: 317-318.