

Balance Outcome Measures – Selection and Utilization

References

- Abrams D, Davidson M, Harrick J, Harcourt P, Zylinski M, Clancy J. Man Ther. 20067.
- Bell DR, Guskiewicz KM, Clark MA, Padua DA. Systematic review of the Balance Error Scoring System. *Sports Health*. 2011. 3: 287-95.
- Cohen ET, Potter K, Allen DD, Bennett SE, Brandfass KG, Widener GL, Yorke AM. Selecting rehabilitation outcome measures for people with multiple sclerosis. *Int J MS Care*. 2015; 17: 181-89.
- Ghislieri M, Gastaldi L, Pastorelli S, Tadano S, Agostini V. Wearable inertial sensors to assess standing balance: A systematic review. *Sensors (Basel)*. 2019; 19(19). pii: E4075. doi: 10.3390/s19194075
- Gribble PA, Hertel J, Plisky P. Using the Star Excursion Balance Test to assess dynamic postural-control deficits and outcomes in lower extremity injury: A literature and systematic review. *J Athletic Training*. 2012; 47: 339-5
- Haines T, Kuys SS, Morrison G, Clarke J, Bew P, McPhail S. Development and validation of the balance outcome measure for elder rehabilitation. *Arch Phys Med Rehabil*. 2007; 88: 1614-21.
- Jette DU, Halbert J, Iverson C, Miceli E, Shah P. Use of standardized outcome measures in physical therapist practice: Perceptions and applications. *Phys Ther*. 2009; 89: 15-35.
- Landers MR, Oscar S, Sasaoka J, Vaughn K. Balance confidence and fear of falling avoidance behavior are most predictive of falling in older adults: Prospective Analysis. *Phys Ther*. 2016; 96: 433-42.
- Lusardi MM, Fritz S, Middleton A et al. Determining risk of falls in community dwelling older adults: A systematic review and meta-analysis using posttest probability. *J Geri Phys Ther*. 2017; 40: 1-36.
- Moore JL, Potter K, Blankshain K, Kaplan SL, O'Dwyer LC, Sullivan Jane E. A core set of outcome measures for adults with neurologic conditions undergoing rehabilitation. *JNPT*. 2018; 42: 174-220.
- Potter K, Fulk GD, Salem Y, Sullivan J. Outcome measures in neurological physical therapy practice: part I. Making sound decisions. *JNPT*. 2011; 35: 57-64.
- Salter KL, Teasell RW, Foley NC, Jutai JW. Outcome assessment in randomized controlled trials of stroke rehabilitation. *Am J Phys Med Rehabil*. 2007; 86: 1007-12.

Sibley KM, Beauchamp MK, Van Ooteghem K, Straus SE, Jaglal SB. Using the systems framework for postural control to analyze the components of balance evaluated in standardized balance measures: A scoping review. *Arch Phys Med Rehabil*. 2015; 96: 122-32.

Tyson SF. How to measure balance in clinical practice. A systematic review of the psychometrics and clinical utility of measures of balance activity for neurological conditions. 2009; 23: 824-40.

Winser SJ, Smith C, Hale LA, Claydon LS, Whitney SL. Balance outcome measures in cerebellar ataxia: a Delphi survey. *Disabil Rehabil*. 2015; 37: 165-70.

Baccini M, Paci M, Rinaldi L. The scale for contraversive pushing: A reliability and validity study. *Neurorehabil Neural Rep*. 2006; 20: 468-72.

Baccini M, Paci M, Nannetti L, Birocolti C, Rinaldi L. Scale for contraversive pushing: cutoff scores for diagnosing pusher behavior and construct validity. *Phys Ther*. 2008; 88: 947-55.

D'Aquila MA, Smith T, Organ D et al. Validation of a lateropulsion scale for patients recovering from stroke. *Clin Rehabil*. 2004; 18: 102-109.

Bergmann J, Krewer C, Riess K, Muller F, Koenig E, Jahn K. Inconsistent classification of pusher behavior in stroke patients: a direct comparison of the Scale for Contraversive Pushing and the Burke Lateropulsion Scale. *Clin Rehabil*. 2014; 28: 696-703.

Babyar SR, Peterson MGE, Bohannon R, Perennou D, Reding M. Clinical examination tools for lateropulsion or pusher syndrome following stroke: a systematic review of the literature. *Clin Rehabil*. 2009; 23: 639-50.

Koter R, Regan S, Clark C, Huang V, Mosley M, Wyant E, Cook C, Hoder J. Clinical outcome measures for lateropulsion poststroke: An updated systematic review. *JNPT*. 2017; 41: 145-55.