

REFERENCES

1. A Hamid MS, Mohamed Ali MR, Yusof A, et al. Platelet-rich plasma injections for the treatment of hamstring injuries: a randomized controlled trial. *Am J Sports Med* 2014;42:2410–18.
2. Askling CM, Tengvar M, Tarassova O, et al. Acute hamstring injuries in Swedish elite sprinters and jumpers: a prospective randomised controlled clinical trial comparing two rehabilitation protocols. *Br J Sports Med* 2014;48:532–9. doi:10.1136/bjsports-2013-093214
3. Askling CM, Nilsson J, Thorstensson A. A new hamstring test to complement the common clinical examination before return to sport after injury. *Knee Surg Sports Traumatol Arthrosc* 2010;18:1798–803
4. Ardern CL, Glasgow P, Schneiders A, Witvrouw E, Clarsen B, Cools A, Gojanovic B, Griffin S, Khan KM, Moksnes H, Mutch SA, Phillips N, Reurink G, Sadler R, Silbernagel KG, Thorborg K, Wangensteen A, Wilk KE, Bizzini M. 2016 Consensus statement on return to sport from the First World Congress in Sports Physical Therapy, Bern. *Br J Sports Med*. 2016 Jul;50(14):853-64. doi: 10.1136/bjsports-2016-096278. Epub 2016 May 25.
5. Ardern CL, Bizzini M, Bahr R. It is time for consensus on return to play after injury: five key questions. *Br J Sports Med* 2016;50:506–8. doi:10.1136/bjsports-2015-095475
6. Ardern CL, Taylor NF, Feller JA, Whitehead TS, Webster KE. Psychological Responses Matter in Returning to Preinjury Level of Sport After Anterior Cruciate Ligament Reconstruction Surgery. *Am J Sports Med*. 2013;41(7):1549-58.
7. Ardern CL. Anterior cruciate ligament reconstruction-not exactly a one-way ticket back to the preinjury level: a review of contextual factors affecting return to sport after surgery. *Sports Health* 2015;7:224–30
8. Ardern CL, Taylor NF, Feller JA, et al. A systematic review of the psychological factors associated with returning to sport following injury. *Br J Sports Med* 2013;47:1120–6.
9. Ardern CL, Österberg A, Tagesson S, et al. The impact of psychological readiness to return to sport and recreational activities after anterior cruciate ligament reconstruction. *Br J Sports Med* 2014;48:1613–19.
10. Ardern CL, Taylor NF, Feller JA, et al. Fifty-five per cent return to competitive sport following anterior cruciate ligament reconstruction surgery: an updated systematic review and meta-analysis including aspects of physical functioning and contextual factors. *Br J Sports Med* 2014;48:1543–52
11. Atkins E, Colville G, John M. A 'biopsychosocial' model for recovery: a grounded theory study of families' journeys after a Paediatric Intensive Care admission. *Intensive Crit Care Nurs* 2012;28:133–40.
12. Askling CM, Tengvar M, Saartok T, et al. Acute first-time hamstring strains during high-speed running: a longitudinal study including clinical and magnetic resonance imaging findings. *Am J Sports Med* 2007;35:19
13. Aune KT, Andrews JR, Dugas JR, et al. Return to play after partial lateral meniscectomy in National Football League athletes. *Am J Sports Med* 2014;42:1865–72
14. Ayers DC, Franklin PD, Ring DC. The role of emotional health in functional outcomes after orthopaedic surgery: extending the biopsychosocial model to orthopaedics. *J Bone Joint Surg Am* 2013;95:e165.
15. Bahr R. Why screening tests to predict injury do not work—and probably never will...: a critical review. *Br J Sports Med*. 2016.
16. Beischer S, Gustavsson L, Senorski EH, Karlsson J, Thomeé C, Samuelsson K, Thomeé R. Young Athletes Who Return to Sport Before 9 Months After Anterior Cruciate Ligament Reconstruction Have a Rate of New Injury 7 Times That of Those Who Delay Return. *J Orthop Sports Phys Ther*. 2020 Feb;50(2):83-90. doi: 10.2519/jospt.2020.9071.
17. Blanch P, Gabbett TJ. Has the athlete trained enough to return to play safely? The acute:chronic workload ratio permits clinicians to quantify a player's risk of subsequent injury. *Br J Sports Med* 2016;50:471–5
18. Brooks JHM. Incidence, risk, and prevention of hamstring muscle injuries in professional rugby union. *Am J Sports Med* 2006;34:1297–306. doi:10.1177/0363546505286022Abstract/FREE Full TextGoogle Scholar
19. Brown R, Orchard J, Kington M, et al. Aprotinin in the management of Achilles tendinopathy: a randomised controlled trial. *Br J Sports Med* 2006;40:275–9. doi:10.1136/bjsm.2005.021931
20. Buckthorpe M, Della Villa F, Della Villa S, Roi GS. On-field Rehabilitation Part 1: 4 Pillars of High-Quality On-field Rehabilitation Are Restoring Movement Quality, Physical Conditioning, Restoring Sport-Specific Skills, and Progressively Developing Chronic Training Load. *J Orthop Sports Phys Ther*. 2019 Aug;49(8):565-569. doi: 10.2519/jospt.2019.8954. Epub 2019 Jul 10.

21. [Buckthorpe M, Della Villa F, Della Villa S, Roi GS. On-field Rehabilitation Part 2: A 5-Stage Program for the Soccer Player Focused on Linear Movements, Multidirectional Movements, Soccer-Specific Skills, Soccer-Specific Movements, and Modified Practice. J Orthop Sports Phys Ther. 2019 Aug;49\(8\):570-575. doi: 10.2519/jospt.2019.8952. Epub 2019 Jul 10.](#)
22. Birmingham P, Muller M, Wickiewicz T, et al. Functional outcome after repair of proximal hamstring avulsions. *J Bone Joint Surg Am* 2011;93: 1819–26. doi:10.2106/JBJS.J.01372Abstract/FREE Full TextGoogle Scholar
23. Cohen SB, Towers JD, Zoga A, et al. Hamstring injuries in professional football players: magnetic resonance imaging correlation with return to play. *Sports Health* 2011;3:423–30
24. Caine DJ, Harmer PA, Schiff MA. Epidemiology of injury in Olympic sports, volume XVI. Oxford, UK: Wiley-Blackwell, 2009.
25. Casartelli NC, Leunig M, Maffiuletti NA, et al. Return to sport after hip surgery for femoroacetabular impingement: a systematic review. *Br J Sports Med* 2015;49:819–24. doi:10.1136/bjsports-2014-094414Abstract/FREE Full Text
26. Clarsen B, Bahr R, Heymans MW, et al. The prevalence and impact of overuse injuries in five Norwegian sports: application of a new surveillance method. *Scand J Med Sci Sport* 2015;25:323–30.
27. Clarsen B, Bahr R, Andersson SH, et al. Reduced glenohumeral rotation, external rotation weakness and scapular dyskinesis are risk factors for shoulder injuries among elite male handball players: a prospective cohort study. *Br J Sports Med* 2014;48:1327–33. doi:10.1136/bjsports-2014-093702
28. Cools AM, Vanderstuyken F, Vereecken F, et al. Eccentric and isometric shoulder rotator cuff strength testing using a hand-held dynamometer: reference values for overhead athletes. *Knee Surg Sports Traumatol Arthrosc* 2015
29. Czuppon S, Racette BA, Klein SE, et al. Variables associated with return to sport following anterior cruciate ligament reconstruction: a systematic review. *Br J Sports Med* 2014;48:356–64.
30. Ellenbecker TS, Cools A. Rehabilitation of shoulder impingement syndrome and rotator cuff injuries: an evidence-based review. *Br J Sports Med* 2010;44:319–27. doi:10.1136/bjsm.2009.058875
31. Elwyn G, Frosch D, Thomson R, et al. Shared decision making: a model for clinical practice. *J Gen Intern Med* 2012;27:1361–7. doi:10.1007/s11606-012-2077-6
32. Ekstrand J, Hägglund M, Waldén M. Epidemiology of muscle injuries in professional football (soccer). *Am J Sports Med* 2011;39:1226–32. doi:10.1177/0363546510395879
33. Ekstrand J, Healy JC, Waldén M, et al. Hamstring muscle injuries in professional football: the correlation of MRI findings with return to play. *Br J Sports Med* 2012;46:112–17. doi:10.1136/bjsports-2011-090155Abstract/FREE Full TextGoogle Scholar
34. Emery CA, Meeuwisse WH, Powell JW. Groin and abdominal strain injuries in The National Hockey League. *Clin J Sport Med* 1999;9:151–6. doi:10.1097/00042752-199907000-00006
35. Engebretsen AH, Myklebust G, Holme I, et al. Intrinsic risk factors for groin injuries among male soccer players: a prospective cohort study. *Am J Sports Med* 2010;38:2051–7. doi:10.1177/0363546510375544
36. Fournier-Farley C, Lamontagne M, Gendron P, et al. Determinants of return to play after the nonoperative management of hamstring injuries in athletes: a systematic review. *Am J Sports Med* 2015; doi:10.1177/0363546515617472. doi: 10.1177/0363546515617472
37. Gabbett TJ, Hulin BT, Blanch P, et al. High training workloads alone do not cause sports injuries: how you get there is the key issue. *Br J Sports Med* 2016;50:444–5.
38. Gabbett TJ, Benton D. Reactive agility of rugby league players. *J Sci Med Sport* 2009;12:212–14.
39. Glazer DD. Development and preliminary validation of the Injury-Psychological Readiness to Return to Sport (I-PRRS) Scale. *J Athl Train* 2009;44:185–9
40. Grindem H, Risberg MA, Eitzen I. Two factors that may underpin outstanding outcomes after ACL reconstruction. *Br J Sports Med* 2015;49:1425
41. Grindem H, Eitzen I, Engebretsen L, et al. Nonsurgical or surgical treatment of ACL injuries: knee function, sports participation, and knee reinjury: the Delaware-Oslo ACL Cohort Study. *J Bone Joint Surg Am* 2014;96:1233–41. doi:10.2106/JBJS.M.01054
42. Gajhede-Knudsen M, Ekstrand J, Magnusson H, et al. Recurrence of Achilles tendon injuries in elite male football players is more common after early return to play: an 11-year follow-up of the UEFA Champions League injury study. *Br J Sports Med* 2013;47:763–8. doi:10.1136/bjsports-2013-092271
43. Hallén A, Ekstrand J. Return to play following muscle injuries in professional footballers. *J Sport Sci* 2014;32:1229–36. doi: 10.1080/02640414.2014.905695CrossRefGoogle Scholar

44. Hägglund M, Waldén M, Ekstrand J. Lower reinjury rate with a coach-controlled rehabilitation program in amateur male soccer: a randomized controlled trial. *Am J Sports Med* 2007;35:1433–42. doi:10.1177/0363546507300063Abstract/FREE Full Text
45. Hägglund M, Waldén M, Ekstrand J. Lower reinjury rate with a coach-controlled rehabilitation program in amateur male soccer: a randomized controlled trial. *Am J Sports Med* 2007;35:1433–42
46. Houshian S, Tscherning T, Riegels-Nielsen P. The epidemiology of Achilles tendon rupture in a Danish county. *Injury* 1998;29:651–4. doi:10.1016/S0020-1383(98)00147-8
47. Hölmich P, Thorborg K, Dehlendorff C, et al. Incidence and clinical presentation of groin injuries in sub-elite male soccer. *Br J Sports Med* 2014;48:1245–50
48. Kvist J, Kartus J, Karlsson J, et al. Results from the Swedish National Anterior Cruciate Ligament Register. *Arthroscopy* 2014;30:803–10.
49. Laprade RF, Surowiec RK, Sochanska AN, et al. Epidemiology, identification, treatment and return to play of musculoskeletal-based ice hockey injuries. *Br J Sports Med* 2014;48:4–10
50. Laprade RF, Wijdicks CA. The management of injuries to the medial side of the knee. *J Orthop Sports Phys Ther* 2012;42:221–33.
51. Lockie RG, Jeffriess MD, McGann TS, et al. Planned and reactive agility performance in semiprofessional and amateur basketball players. *Int J Sports Physiol Perform* 2014;9:766–71
52. Laudner KG, Stanek JM, Meister K. Differences in scapular upward rotation between baseball pitchers and position players. *Am J Sports Med* 2007;35:2091–5
53. Lorenz D, Domzalski S. CRITERIA-BASED RETURN TO SPRINTING PROGRESSION FOLLOWING LOWER EXTREMITY INJURY. *Int J Sports Phys Ther*. 2020 Apr;15(2):326-332.
54. Losciale JM, Zdeb RM, Ledbetter L, Reiman MP, Sell TC. The association between passing return-to-sport criteria and second anterior cruciate ligament injury risk: a systematic review with meta-analysis. *J Orthop Sports Phys Ther*. 2018;49(2):43-54.
55. Lundblad M, Waldén M, Magnusson H, et al. The UEFA injury study: 11-year data concerning 346 MCL injuries and time to return to play. *Br J Sports Med* 2013;47:759–62.
56. Kilcoyne KG, Dickens JF, Keblish D, et al. Outcome of grade I and II hamstring injuries in intercollegiate athletes: a novel rehabilitation protocol. *Sports Health* 2011;3:528–33. doi:10.1177/1941738111422044Abstract/FREE Full TextGoogle Scholar
57. King E, Ward J, Small L, et al. Athletic groin pain: a systematic review and meta-analysis of surgical versus physical therapy rehabilitation outcomes. *Br J Sports Med* 2015;49:1447–51. doi:10.1136/bjsports-2014-093715Abstract/FREE Full TextGoogle Scholar
58. Kujala UM, Sarna S, Kaprio J. Cumulative incidence of Achilles tendon rupture and tendinopathy in male former elite athletes. *Clin J Sports Med* 2005;15:133–5.
59. Jacobsen P, Witvrouw E, Muxart P, et al. A combination of initial and follow-up physiotherapist examination predicts physician-determined time to return to play after hamstring injury, with no added value of MRI. *Br J Sports Med* 2016;50:431–9. doi:10.1136/bjsports-2015-095073
60. Moen MH, Reurink G, Weir A, et al. Predicting return to play after hamstring injuries. *Br J Sports Med* 2014;48:1358–63.
61. Malliaropoulos N, Papacostas E, Kiritsi O, et al. Posterior thigh muscle injuries in elite track and field athletes. *Am J Sports Med* 2010;38:1813–19. doi:10.1177/0363546510366423
62. Maletis GB, Inacio MC, Funahashi TT. Risk factors associated with revision and contralateral anterior cruciate ligament reconstructions in the Kaiser Permanente ACLR Registry. *Am J Sports Med* 2015;43:641–7.
63. Myklebust G, Bahr R. Return to play guidelines after anterior cruciate ligament surgery. *Br J Sports Med* 2005;39:127–31.
64. Möller M, Movin T, Granhed H, et al. Acute rupture of tendon Achilles. A prospective randomised study of comparison between surgical and non-surgical treatment. *J Bone Joint Surg Br* 2001;83:843–8
65. Mullaney MJ, McHugh MP, Tyler TF, et al. Weakness in end-range plantar flexion after Achilles tendon repair. *Am J Sports Med* 2006;34:1120–5. doi:10.1177/0363546505284186Abstract/FREE Full TextGoogle Scholar
66. Nawabi DH, Cro S, Hamid IP, et al. Return to play after lateral meniscectomy compared with medial meniscectomy in elite professional soccer players. *Am J Sports Med* 2014;42:2193–8
67. Orchard JW, Seward H, Orchard JJ. Results of 2 decades of injury surveillance and public release of data in The Australian Football League. *Am J Sports Med* 2013;41:734–41

68. Orchard J, Seward H. Epidemiology of injuries in The Australian Football League, seasons 1997–2000. *Br J Sports Med* 2002;36:39–44. doi:10.1136/bjsm.36.1.39Abstract/FREE Full TextGoogle Scholar
69. Parekh SG, Wray WH, Brimmo O, et al. Epidemiology and outcomes of Achilles tendon ruptures in The National Football League. *Foot Ankle Spec* 2009;2:283–6
70. Paterno MV, Schmitt LC, Thomas S, Duke N, Russo R, Quatman-Yates CC.. Patient and Parent Perceptions of Rehabilitation Factors That Influence Outcomes After Anterior Cruciate Ligament Reconstruction and Clearance to Return to Sport in Adolescents and Young Adults. *J Orthop Sports Phys Ther*. 2019 Aug;49(8):576-583. doi: 10.2519/jospt.2019.8608. Epub 2019 Feb 13.
71. Pincus T, Kent P, Bronfort G, et al. Twenty-five years with the biopsychosocial model of low back pain—is it time to celebrate? A report from the Twelfth International Forum for Primary Care Research on Low Back Pain. *Spine* 2013;38:2118–23.
72. Podlog L, Dimmock J, Miller J. A review of return to sport concerns following injury rehabilitation: Practitioner strategies for enhancing recovery outcomes. *Phys Ther Sport*. 2011;12(1):36-42.
73. Pollock N, Patel A, Chakraverty J, et al. Time to return to full training is delayed and recurrence rate is higher in intratendinous ('c') acute hamstring injury in elite track and field athletes: clinical implication of the British Athletics Muscle Injury Classification. *Br J Sports Med* 2016;50:305–10
74. Reurink G, Brilman EG, de Vos RJ, et al. Magnetic resonance imaging in acute hamstring injury: can we provide a return to play prognosis? *Sports Med* 2015;45:133–46. doi:10.1007/s40279-014-0243-1
75. Ribeiro A, Pascoal AG. Resting scapular posture in healthy overhead throwing athletes. *Man Ther* 2013;18:547–50.
76. Røtterud JH, Sivertsen EA, Forssblad M, et al. Effect of meniscal and focal cartilage lesions on patient-reported outcome after anterior cruciate ligament reconstruction: a nationwide cohort study from Norway and Sweden of 8476 patients with 2-year follow-up. *Am J Sports Med* 2013;41:535–43
77. Sanfilippo JL, Silder A, Sherry MA, et al. Hamstring strength and morphology progression after return to sport from injury. *Med Sci Sports Exerc* 2013;45:448–54. doi:10.1249/MSS.0b013e3182776eff
78. Silder A, Sherry MA, Sanfilippo J, et al. Clinical and morphological changes following 2 rehabilitation programs for acute hamstring strain injuries: a randomized clinical trial. *J Orthop Sports Phys Ther* 2013;43:284–99
79. Smith AH, Capin J, Zarzycki R, Snyder-Mackler LAthletes With Bone-Patellar Tendon Bone Autograft for ACL Reconstruction Were Months Slower to Meet Rehabilitation Milestones and Return to Sport Criteria Than Athletes With Hamstring Tendon Autograft or Soft Tissue Allograft: Secondary Analysis From the ACL-SPORTS Trial. *J Orthop Sports Phys Ther*. 2019 Nov 27:1-28. doi: 10.2519/jospt.2020.9111. [Epub ahead of print]
80. Serner A, van Eijck CH, Beumer BR, et al. Study quality on groin injury management remains low: a systematic review on treatment of groin pain in athletes. *Br J Sports Med* 2015;49:813
81. Shitara H, Kobayashi T, Yamamoto A, et al. Prospective multifactorial analysis of preseason risk factors for shoulder and elbow injuries in high school baseball pitchers. *Knee Surg Sports Traumatol Arthrosc* 2015
82. Shrier I, Safai P, Charland L. Return-to-play following injury: whose decision should it be? *Br J Sports Med* 2014;48:394–401.
83. Shrier I. Strategic Assessment of Risk and Risk Tolerance (StARRT) framework for return-to-play decision-making. *Br J Sports Med* 2015;49:1311–15.
84. Sheppard JM, Young WB, Doyle TLA, et al. An evaluation of a new test of reactive agility and its relationship to sprint speed and change of direction speed. *J Sci Med Sport* 2006;9:342–9
85. Silbernagel KG, Thomeé R, Thomeé P, et al. Eccentric overload training for patients with chronic Achilles tendon pain—a randomised controlled study with reliability testing of the evaluation methods. *Scand J Med Sci Sport* 2001;11:197–206. doi: 10.1034/j.1600-0838.2001.110402.x
86. Silbernagel KG, Thomeé R, Eriksson BI, et al. Continued sports activity, using a pain-monitoring model, during rehabilitation in athletes with Achilles tendinopathy: a randomized controlled study. *Am J Sports Med* 2007;35:897–906.
87. Silbernagel KG, Steele R, Manal K. Deficits in heel-rise height and Achilles tendon elongation occur in patients recovering from an Achilles tendon rupture. *Am J Sports Med* 2012;40:1564–71
88. Silbernagel KG, Crossley KM. A proposed return to sport program for patients with midportion Achilles tendinopathy: rationale and implementation. *J Orthop Sports Phys Ther* 2015;21:1–42
89. Sherry MA, Best TM. A comparison of 2 rehabilitation programs in the treatment of acute hamstring strains. *J Orthop Sports Phys Ther* 2004;34:116–25

90. Tol JL, Hamilton B, Eirale C, et al. At return to play following hamstring injury the majority of professional football players have residual isokinetic deficits. *Br J Sports Med* 2014;48:1364–9
91. Thorborg K, Branci S, Nielsen M, et al. Eccentric and isometric hip adduction strength in male soccer players with and without adductor-related groin pain. An assessor-blinded comparison. *Orthop J Sports Med* 2014;2:2325967114521778. [Abstract/FREE Full TextGoogle Scholar](#)
92. Thorborg K, Hölmich P. Advancing hip and groin injury management: from eminence to evidence. *Br J Sports Med* 2013;47:602–5. doi:10.1136/bjsports-2012-092090FREE Full TextGoogle Scholar
93. Tripp DA, Stanish W, Ebel-Lam A, et al. Fear of reinjury, negative affect, and catastrophizing predicting return to sport in recreational athletes with anterior cruciate ligament injuries at 1 year postsurgery. *Rehabil Psych* 2007;52:74–81.
94. Timmins RG, Bourne MN, Shield AJ, et al. Short biceps femoris fascicles and eccentric knee flexor weakness increase the risk of hamstring injury in elite football (soccer): a prospective cohort study. *Br J Sports Med* Published Online First: 16 Dec 2015 doi: 10.1136/bjsports-2015-095362
95. Tyler TF, Nicholas SJ, Campbell RJ, et al. The association of hip strength and flexibility with the incidence of adductor muscle strains in professional ice hockey players. *Am J Sports Med* 2001;29:124–8.
96. Wangensteen A, Almusa E, Boukarroum S, et al. MRI does not add value over and above patient history and clinical examination in predicting time to return to sport after acute hamstring injuries: a prospective cohort of 180 Male athletes. *Br J Sports Med* 2015;49:1579–87. doi:10.1136/bjsports-2015-094892
97. Webster KE, Feller JA, Leigh WB, et al. Younger patients are at increased risk for graft rupture and contralateral injury after anterior cruciate ligament reconstruction. *Am J Sports Med* 2014;42:641–
98. Webster KE, Feller JA. Expectations for return to preinjury sport before and after anterior cruciate ligament reconstruction. *Am J Sports Med*. 2019;47(3):578-83.
99. Webster KE, Feller JA, Lambros C. Development and preliminary validation of a scale to measure the psychological impact of returning to sport following anterior cruciate ligament reconstruction surgery. *Phys Ther Sport* 2008;9:9–15.
100. Werner J, Hägglund M, Walden M, et al. UEFA injury study: a prospective study of hip and groin injuries in professional football over seven consecutive seasons. *Br J Sports Med* 2009;43:1036–40.
101. Wilk KE, Macrina LC, Fleisig GS, et al. Deficits in glenohumeral passive range of motion increase risk of shoulder injury in professional baseball pitchers: a prospective study. *Am J Sports Med* 2015;43:2379–85. doi:10.1177/0363546515594380
102. Wiese-Bjornstal DM, Smith AM, Shaffer SM, et al. An integrated model of response to sport injury: psychological and sociological dynamics. *J Appl Sport Psychol* 1998;10:46–69.
103. Windt J, Gabbett TJ. How do training and competition workloads relate to injury? The workload–injury aetiology model. *Br J Sports Med*. 2017;51(5):428.
104. Wright RW, Huston LJ, Spindler KP, et al., MOON Group. Descriptive epidemiology of the Multicenter ACL Revision Study (MARS) cohort. *Am J Sports Med* 2010;38:1979–86
105. Yeung SS, Suen AMY, Yeung EW. A prospective cohort study of hamstring injuries in competitive sprinters: preseason muscle imbalance as a possible risk factor. *Br J Sports Med* 2009;43:589–94

CVI.