

## Walking Recovery Post Stroke References

Abbasian S, Rastegar M. Is the intensity or duration of treadmill training important for stroke patients? A meta-analysis. *J Stroke Cerebrovasc Dis*. 2018; 27: 32-43.

Holleran CL, Straube DD, Kinnaird CR et al. Feasibility and potential efficacy of high intensity stepping training in variable contexts in subacute and chronic stroke. *Neurorehabil Neural Rep*. 2014; 28: 643-51.

Hornby TG, Holleran CL, Hennessy PW et al. Variable intensive early walking poststroke (VIEWS): A randomized controlled trial. *Neurorehabil Neural Rep*. 2016; 30: 440-50.

Hornby TG, Reisman DS, Ward IG et al. Clinical Practice Guideline to improve locomotor function following chronic stroke, incomplete spinal cord injury and brain injury. *J Neurol Phys Ther*. 2020; 44: DOI: 10.1097/NPT.0000000000000303

Wiener J, McIntyre A, Janssen S, Chow JT, Batey C, Teasell R. Effectiveness of high-intensity interval training for fitness and mobility post stroke: A systematic review. *PM R*. 2019; epub ahead of print. Doi: 10.1002/pmrj.12154.

Williams G, Hassett L, Clark R, Bryant A, Olver J, Morris ME, Ada L. Improving walking ability in people with neurologic conditions: A theoretical framework for biomechanics-driven exercise prescription. *Arch Phys Med Rehabil*. 2019; 100: 1184-90.

Winstein CJ et al. Guidelines for adult stroke rehabilitation and recovery. A guideline for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke*. 2016. 47: e98e-169.